



LEGEND is a broad spectrum systemic fungicide for winter and spring crops of wheat (excluding durum), barley, oats, winter rye, oilseed rape, field beans, cabbages, carrots, leeks, parsnips, swedes, turnips and horseradish.

An emulsion (oil in water) formulation containing 250 g/litre tebuconazole.

SAFETY PRECAUTIONS

Operator Protection

WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.
WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.
WHEN USING DO NOT EAT, DRINK OR SMOKE.
IN CASE OF CONTACT WITH EYES RINSE IMMEDIATELY with plenty of water and seek medical advice.
WASH HANDS AND EXPOSED SKIN before meals and after work.
IF YOU FEEL UNWELL, seek medical advice (show label where possible).

Environmental Protection

Do not contaminate ponds, waterways or ditches with the product or its container. (Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads).
Aim spray away from water.

Storage and Disposal

DO NOT RE-USE CONTAINER FOR ANY PURPOSE.
KEEP IN ORIGINAL CONTAINER tightly closed in a safe place.
KEEP OUT OF REACH OF CHILDREN.
KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS.
WASH OUT CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

Net contents
4 x 5L e

Authorisation Holder and Marketing Company
JT Agro Europe sp. z o.o.
Gate A, Aleja Grunwaldza 472, 80-309 Gdańsk, Poland
Telephone: +44 1484 848885
www.jtcrop.com
info@jtcrop.com



GROUP 3 FUNGICIDE

In the event of an emergency, call the National Poisons Information Centre,
Beaumont Hospital: **01-8092166** or **01-8379964**

LEGEND Contains 250 g/L (25.9%) tebuconazole.	SAFETY INFORMATION	UFI: 1VV2-T0A1-Q00X-YCQ0
 DANGER Harmful if swallowed. Causes serious eye damage. Harmful if inhaled. Suspected of damaging the unborn child. Very toxic to aquatic life with long lasting effects. <i>Refer to the leaflet attached to the product pack for full Safety Information.</i>		
		PCS 06472

IMPORTANT INFORMATION				
FOR USE ONLY AS AN AGRICULTURAL/HORTICULTURAL FUNGICIDE				
Crop	Maximum single dose	Maximum number of applications	Maximum total dose	Latest time of application
Winter wheat, spring wheat, winter barley, spring barley, oats, winter rye	1 L/ha	–	1 L/ha	Before grain watery ripe stage (GS 71)
Oilseed rape (winter & spring)	1 L/ha	–	2 L/ha	End of flowering (BBBH 69)
Cabbage, field bean, linseed, swede, turnip	1 L/ha	–	1 L/ha	35 days before harvest
Carrot, horseradish, parsnip	1 L/ha	–	1 L/ha	21 days before harvest
Leek	1 L/ha	–	1 L/ha	14 days before harvest

Other specific restrictions:
To protect aquatic organisms respect an unsprayed buffer zone of 5 M to surface water bodies.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

**PROTECT FROM FROST
SHAKE WELL BEFORE USE**